

The Ordovician System in Estonia

Tõnu Meidla, Leho Ainsaar and Olle Hints

The following text is based on Meidla et al. (2023).

The area of continuous distribution of the Ordovician in the Baltic Sea area and east of it extends from the southern part of the Baltic Sea in the west to the vicinity of Moscow in the east and from the Gulf of Finland or northernmost Estonia in the north to Belarus and Poland in the south. In the northern part of this area, in the eastern coastal region of the Baltic Sea, beds are exposed in sections of the Baltic-Ladoga Klint, sometimes with more than 10 m of strata exposed, which attracted the attention of early investigators, together with other coastal and river bank sections, as well as old and new limestone quarries and open cast pits. Well-accessible successions with well-preserved fossils and sedimentary structures attracted the attention of investigators already in the early 19th century and were addressed in the papers by O. M. L. v. Engelhardt (1820), W. T. H. F. Strangways (1821), E. Eichwald (1825), and others. In particular, the characteristic succession of the Cambrian to Middle Ordovician with a number of distinctive rock units (the Cambrian Blue Clay, phosphatic brachiopod coquina, *Dictyonema* argillite, dark green glauconite sandstone and the succession of distinctive limestone units above) was of great interest.

The thorough monographic paper by F. Schmidt from 1858 brought to attention the main features of Ordovician stratigraphy in Estonia for the first time. The general pattern of his very old geologic map in the same volume is already well demonstrating the most characteristic feature of modern bedrock maps of Estonia —latitudinally orientated outcrop belts of the Ordovician and Silurian stages in northern Estonia. The feature is a result of the generally simple geologic structure of the area, with almost horizontal strata dipping southward only 2–5 m/km.

The Lower Ordovician thin succession of siliciclastics comprises sandstones, argillites and clays (Pakerort and Varangu RSs) overlain by the glauconitic sand- and siltstones (Hunneberg and Billingen RSs). Further up, the main part of the Ordovician succession in northern Estonia is heavily dominated by various kinds of limestones but also contains some intercalations of kukersite oil shale concentrated mainly in the Kukruse Regional Stage (RS). The transition from the terrigenous to carbonate rocks in the basal part of the Toila Formation (topmost Lower Ordovician to basal Middle Ordovician) is marked by the fairly sharp appearance of the first limestone/dolomite beds. The appearance of the first representatives of the numerous characteristic Middle Ordovician fossil groups is recorded in the same transition interval or in the overlying Volkhov Stage.

The Ordovician limestone succession in Estonia and ad-

jacent areas begins with carbonates deposited in a sediment-starved shallow marine basin. Upward the sedimentation rates have increased in obvious correlation with the carbonate production. In the Upper Ordovician, corals make their first appearance, and the first carbonate buildups appear, emphasising a remarkable change in the overall character of the palaeobasin. Informer publications, change in the type of sedimentation and the character of biofacies is ascribed to a gradual climatic change resulting from the northward drift of the Baltica Palaeocontinent from the temperate climatic zone to the (sub)tropical realm (Nestor and Einasto, 1997). This interpretation, however, is not in full accordance with the results of a pilot study on conodont phosphate oxygen isotope palaeotemperature revealing a continuous cooling trend in the palaeobasin throughout the Middle and Later Ordovician (Männik et al. 2022). Independently of that, the Middle and Upper Ordovician changes resulted in an increase of carbonate production and sedimentation rate on the carbonate shelf where the deposition pattern was obviously controlled by accommodation space available.

The problems of the Ordovician geology in the subsurface area in central and southern Estonia were first revealed in the late 1950s when the extensive drilling started in the area. Thanks to the high number of drill cores obtained during a comprehensive drilling programme in the 1950s–1980s, the main correlations problems between the stratigraphic successions in the outcrop area and in southern Estonia were mainly resolved. As a result of the comparison of the eastern Baltic and Scandinavian successions, the concepts of the structural-facies zones (by Männil 1966) or confacies belts (by Jaanusson 1976) were introduced for the Ordovician of Baltoscandia (see Fig. 1). As the term “confacies” is unique (being exclusively used for the Ordovician of Baltoscandia only), a different terminology has been introduced by Harris et al. 2004 (see explanation to the Fig. 1). The micropalaeontological and macrofaunal studies of the core sections also revealed the distinctive biogeographic differentiation pattern, characteristic of the Ordovician rocks (Männil 1966; Männil et al. 1968; Meidla 1996, etc.). Although the biofacies pattern is generally described for the eastern Baltic area, the facies zonation of the entire Baltoscandian area is still imperfectly known. The seismic investigations of the Baltic Sea area, performed in the last decades (Tuuling 1998 and references therein), but also detailed (micro)palaeontological investigations (e.g. Tinn 2002) might produce valuable new information in this field.

The total thickness of the Ordovician reaches up to 190 m, being maximal in central and eastern Estonia and considerably less in the outcrop area, as well as in the southwestern mainland of Estonia. Several correlation




Fig. 4. Ordovician stratigraphy of Estonia. Graptolite zonation according to Kaljo & Vingissaar 1969; Kaljo et al. 1986; Männil 1976; Resheniya... 1987; Männil & Meidla 1994; Nölvak et al. 2006; conodont zones according to Männil in Nölvak et al. 2006; Paiste et al. 2020, 2022 and Meidla et al. 2023. Numbers in the column of the conodont zonation correspond to the conodont subzones as follows: subzones of the *Baltoniodus norrandicus* Zone: 1 – *Trapezognathus quadrangulum* Subzone, 2 – *Lenodus antivariabilis* Subzone; subzones of the *Pygodus serra* Zone: 3 – *Eoplacognathus foliaceus* Subzone, 4 – *Eoplacognathus reclinator* Subzone, 5 – *Eoplacognathus robustus* Subzone, 6 – *Eoplacognathus protoamosus* Subzone, 7 – *Eoplacognathus lindstroemi* Subzone. Abbreviations: Llan., Llandovery; Da., Darriwilian; Hir., Hirnantian; Rhu., Rhuddanian; Cyst., Cystograptus; Par., Parakidograptus; Met., Metabolograptus; Dic., Dicellograptus; Nemagr., Nemagraptus; Holmogr., Holmograptus; Phyllogr., Phyllograptus; Rhabdin, Rhabdinopora; flabellif., flabelliforme; Amorphog., Amorphognathus; Baltoniod., Baltoniodus; Yangtzeplac., Yangtzeplacognathus; origin., originalis; trian., triangularis.

problems still persist in the Ordovician of Estonia due to marked biofacies differences between northern and southern Estonia. In part, they are also discussed in a recent monographic overview of Estonian geology (see Heinsalu and Viira 1997, Meidla 1997; Hints 1997; Hints and Meidla 1997 in Raukas and Teedumäe, 1997). New prospects in this field have already been opened by stable isotope studies, as the stable carbon isotope curves have demonstrated a good correlation potential (Kaljo et al. 2004, 2007, and references therein; Ainsaar et al. 2004, 2007, 2010).

The development of the stratigraphic classification of the Ordovician strata in Estonia, from the “beds” (*Schichten*) by Schmidt (1858) to the stages in modern meaning, is documented in detail in Männil (1966), Rõõmusoks (1983) and Rõõmusoks et al. (1997). The term “Ordovician” was likely used to describe the Estonian succession for the first time by Bassler (1911) and became widely established in the geological literature of Estonia since Bekker (1921). A number of regional series and subseries for the Ordovician System in Estonia and neighbouring Russia were introduced by Schmidt (1881) and several subsequent authors. Raymond (1916) introduced the traditional American three-fold subdivision of the Ordovician System for this particular area, but this classification was subjected to repeated changes until 1987. Also, the terms “Oeland Series”, “Viru Series” and “Harju Series” have been widely used as a basic three-fold classification for the Ordovician System of the area since the 1950s (introduced by Kaljo et al. 1958 and Jaanusson 1960 in a nearly recent meaning). The subseries have been introduced as well (see Männil and Meidla 1994 and Nõlvak et al. 2006 for a summary), but they lost their actuality and are rarely used today.

The modern three-fold classification of the Ordovician System (IUGS 2004) was first used for the Estonian succession by Webby (1998) and is presented here in detail (Fig. 4). In relation to the definition of the GSSP for the base of the Ordovician System in the Green Point section, Newfoundland (Remane, 2003), a revision of the traditional position of the Cambrian-Ordovician boundary at the base of the Pakerort RS in Estonia turned out to be necessary. According to conodont data, the system boundary in the northern Estonian sections lies some metres higher than previously suggested, i.e. in the middle of the Pakerort Stage, within the Kallavere Formation (Puura and Viira 1999). The upper boundary (the lower boundary of the Silurian System) has for a long time been correlated with the major hiatus between the Porkuni and Juuru RSs, corresponding to the maximum regression related to the Hirnantian glaciation, due to the fact that a major faunal overturn is recorded on this level. Recent chemostratigraphic correlations, however, suggest that the falling limb of HICE reaches into the basal part of the Juuru RS (Ainsaar et al. 2010, 2015; Bauert et al. 2014; Meidla et al. 2020, 2023) and the Ordovician–Silurian boundary is located within the beds formerly attributed to the basal Silurian.

The term “(Regional) Stage”, first applied by Bekker

(1921), has become the principal category in the chronostratigraphic classification of the Ordovician System in Estonia.

The main features of the chronostratigraphic classification of the Ordovician System were established already by Männil (1966). Only minor changes were introduced in the later decades: the Ceratopyge RS was renamed the Varangu RS (Männil, 1990), the Latorp RS was replaced by the Hunneberg and Billingen RSs (Hints et al., 1993) and a new unit, the Haljala RS, is used instead of the Idavere and Jõhvi RSs (following Jaanusson, 1995 and Nõlvak, 1997). Hints and Nõlvak (1999) brought the concept of boundary stratotypes (“golden spike”) into the Estonian stratigraphy, proposing a stratotype — the Pääsküla outcrop — for the lower boundary of the Keila RS. However, as stratigraphic hiatuses on the stage boundaries are very common in northern Estonia (remarkable faunal changes are usually related to hiatuses), wide usage of this concept for the stage boundaries in this area looks rather complicated.

The lithostratigraphic classification of the Ordovician rocks was introduced by Orviku (1940) for the upper Middle Ordovician. This approach was widely accepted by subsequent authors and led to the compilation of a series of detailed correlation charts approved by the Interdepartmental Stratigraphic Committee of the former USSR (Resheniya... 1965, 1978, 1987 and a related paper by Männil and Rõõmusoks, 1984). The last version of such a formal correlation chart (the edition of 1987) was, in a slightly emended form, published also in English in the series of the IUGS publications (Männil and Meidla, 1994).

The correlation chart in Fig. 4 contains some recent improvements compared to this publication, the most recent ones being introduced by Ainsaar and Meidla (2001), Nõlvak et al. (2006), Meidla et al. (2020 and references therein) and Paiste et al. (2022, 2023). Some more versions of the Ordovician correlation charts for Estonia have also been published by Hints et al. (1993), Nõlvak (1997) and Meidla et al. (2014).

The composition and textures of the Ordovician carbonate rocks and the principal differences between the confacies belts were summarised by Põlma (1982 and references therein).

The monographic studies on Ordovician palaeontology started already in the 19th century. After the comprehensive review of the Ordovician and Silurian strata (in modern meaning) by Schmidt (1858 and several subsequent monographic papers), a number of important monographic papers were published by F. B. Rosen, W. Dybowski, A. Pahlen, G. Holm, A. Mickwitz, O. Jaekel, J. H. Bonnema and R. F. Bassler. The tradition of palaeontological investigations on the Ordovician material of Estonia was continued by A. Öpik (1930, 1934 and others) and, later on, by the recent generation of palaeontologists. Monographs and extensive monographic papers were published on the Ordovician brachiopods, corals,

stromatoporoids, chitinozoans, scolecodonts, ostracods, conodonts, etc. Summaries on the palaeontological investigations on virtually all fossil groups recorded from

the Ordovician of Estonia are published in the recent monograph "Geology and mineral resources of Estonia" (Raukas and Teedumäe 1997).

References

- Ainsaar, L., Meidla, T., 2001. Facies and stratigraphy of the middle Caradoc mixed siliciclastic-carbonate sediments in eastern Baltoscandia. *Proceedings of the Estonian Academy of Sciences. Geology*, **50**, 5–23.
- Ainsaar, L., Meidla, T., Martma, T., 2004. The Middle Caradoc facies and faunal turnover in the Late Ordovician Baltoscandian palaeobasin. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **210**, 119–133.
- Ainsaar, L., Meidla, T., Tinn, O., Martma, T., Dronov, A., 2007. Darriwilian (Middle Ordovician) carbon isotope stratigraphy in Baltoscandia. The Global Ordovician and Silurian. *Acta Palaeontologica Sinica*, **46** (Suppl.), 1–8.
- Ainsaar, L., Kaljo, D., Martma, T., Meidla, T., Männik, P., Nõlvak, J., Tinn, O., 2010. Middle and Upper Ordovician carbon isotope chemostratigraphy in Baltoscandia: A correlation standard and clues to environmental history. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **294**, 189–201.
- Ainsaar, L., Truumees, J., Meidla, T., 2015. The position of the Ordovician–Silurian boundary in Estonia tested by high-resolution $\delta^{13}\text{C}$ chemostratigraphic correlation. In: *Chemostratigraphy: Concepts, Techniques and Applications* (Ramkumar, M., ed.). Amsterdam, Elsevier, p. 395–412.
- Bauert, H., Ainsaar, L., Põldsaar, K., Sepp, S., 2014. $\delta^{13}\text{C}$ chemostratigraphy of the Middle and Upper Ordovician succession in the Tartu-453 drillcore, southern Estonia, and the significance of the HICE. *Estonian Journal of Earth Sciences*, **63**, 195–200.
- Bassler, R. F., 1911. The Early Palaeozoic Bryozoa of the Baltic Provinces. *Smithsonian Institute, US National Museum Bulletin*, **77**, 1–382.
- Bekker, H., 1921. The Kuckers Stage of the Ordovician Rocks of NE Estonia. *Acta et Commentationes Universitatis Tartuensis*, **A2**, 81–84.
- Brenchley, P. J., Carden, G. A., Hints, L., Kaljo, D., Marshall, J.D., Martma, T., Meidla, T., Nõlvak, J., 2003. High-resolution stable isotope stratigraphy of Upper Ordovician sequences: Constraints on the timing of bioevents and environmental changes associated with mass extinction and glaciation. *Geological Society of America, Bulletin*, **115**, 89–104.
- Chen, X., Rong, J.-y., Fan, J., Zhan, R., Mitchell, E., Harper, D. A. T., Melchin, M. J., Peng, P.-a., Finney, S., Wang, X., 2006. The Global Stratotype Section and Point (GSSP) for the base of the Hirnantian Stage (the uppermost of the Ordovician System). *Episodes*, **29**, 183–196.
- Eichwald, E., 1825. *Geognostico-zoologicae per Ingriam marisque Baltici provincias nec non de trilobitis observationes*. Casani, 58 pp.
- Eichwald, E., 1854. Die Grauwackenschichten von Liv- und Estland. *Bull. Soc. Nat. Moscou*, **27**, 1–111.
- Einasto, R., 2007. Ordoviitsiumi ja Siluri ladestu piiri leidmine Neitla kruusakarjääris. *Keskkonnatehnika*, **7**, 60–61.
- Engelhardt, M., 1820. *Darstellung aus dem Felsgebäude Russlands. Erste Lieferung*. Geognostischer Umriss von Finland. Berlin. 44 pp.
- Harris, M. T., Sheehan, P. M., Ainsaar, L., Hints, L., Männik, P., Nõlvak, J., Rubel, M., 2004. Upper Ordovician Baltoscandian sequences of western Estonia. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **210**, 135–148.
- Heinsalu, H., Viira V., 1997. Pakerort Stage. Varangu Stage. In: *Geology and Mineral Resources of Estonia* (Raukas, A., Teedumäe, A., eds). Estonian Academy Publishers, Tallinn, p. 52–58.
- Hints, L., Meidla, T., 1997. Keila Stage. Oandu Stage. Rakvere Stage. Nabala Stage. Vormsi Stage. Pirgu Stage. Porkuni Stage. In: *Geology and Mineral Resources of Estonia* (Raukas, A., Teedumäe, A., eds). Estonian Academy Publishers, Tallinn, p. 74–88.
- Hints, L., 1997. Aseri Stage. Lasnamägi Stage. Uhaku Stage. Haljala Stage. In: *Geology and Mineral Resources of Estonia* (Raukas, A., Teedumäe, A., eds). Estonian Academy Publishers, Tallinn, p. 66–74.
- Hints, L., Meidla, T., Gailite, L.-I., Sarv, L. (comp.), 1993. *Catalogue of Ordovician Stratigraphical Units and Stratotypes of Estonia and Latvia*. Institute of Geology, Estonian Academy of Sciences, Tallinn. 62 pp.
- Hints, L., Oraspõld, A., Kaljo, D., 2000. Stratotype of the Porkuni Stage with comments on the Rõa Member (Uppermost Ordovician, Estonia). *Proceedings of the Estonian Academy of Sciences. Geology*, **49**, 177–199.
- Hints, O., Nõlvak, J., 1999. Proposal for the lower boundary-stratotype of the Keila Regional Stage (Upper Ordovician). *Proceedings of the Estonian Academy of Sciences. Geology*, **48**, 158–169.
- Holland, C. H., 1985. Series and stages of the Silurian System. *Episodes*, **8**, 101–103.
- IUGS, 2004. International stratigraphic chart. IUGS International Commission of Stratigraphy: <http://www.stratigraphy.org/cheu.pdf>
- Jaanusson, V., 1960. Graptoloids from the Ontikan and Viruan (Ordovician) limestones of Estonia and Sweden. *The Bulletin of the Geological Institutions of the University of Uppsala*, **38**, 289–366.
- Jaanusson, V., 1976. Faunal dynamics in the Middle Ordovician (Viruan) of Balto-Scandia. In: *The Ordovician System. Proceedings of a Palaeontological Association Symposium, Birmingham, September 1974* (Bassett, M.G., ed.), University of Wales Press, Cardiff, p. 301–326.
- Jaanusson, V., 1995. Confacies differentiation and upper Middle Ordovician correlation in the Baltoscandian basin. *Proceedings of the Estonian Academy of Sciences. Geology*, **44**, 73–86.
- Kaljo, D., Hints, L., Hints, O., Martma, T., Nõlvak, J., 1999. Carbon isotope excursions and coeval biotic–environmental changes in the late Caradoc and Ashgill of Estonia. *Acta Universitatis Carolinae, Geologica*, **43**, 507–510.
- Kaljo, D., Hints, L., Martma, T. and Nõlvak, J. 2001. Carbon isotope stratigraphy in the latest Ordovician of Estonia. *Chemical Geology*, **175**, 49–59.
- Kaljo, D., Hints, L., Martma, T., Nõlvak, J. and Oraspõld, A. 2004. Late Ordovician carbon isotope trend in Estonia, its significance in stratigraphy and environmental analysis. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **210**, 165–185.
- Kaljo, D., Martma, T., Saadre, T., 2007. Post-Hunnebergian Ordovician carbon isotope trend in Baltoscandia, its environmental implications and some similarities with that of Nevada. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **245**, 138–155.
- Kaljo, D., Martma, T., Männik, P., Viira, V., 2003. Implications of Gondwana glaciations in the Baltic late Ordovician and Silurian and a carbon isotopic test of environmental cyclicity. *Bulletin de la Societe Geologique de France*, **174**, 59–66.
- Kaljo, D., Rõõmusoks, A., Männik, R., 1958. On the series of the Baltic Ordovician and their significance. *Eesti NSV Teaduste Akadeemia Toimetised. Tehniliste ja Füüsikalise-Matemaatiliste Teaduste Seeria*, **7**, 71–74.
- Mägi, S., 1990. Ontika clint. In: *Field meeting Estonia 1990. An Excursion Guidebook* (Kaljo, D., Nestor, H., eds), Institute of Geology, Estonian Academy of Sciences, Tallinn, p. 148–153.
- Mägi, S., 1991. Jägala Fall. In: *Geology and mineral resources of*

- Estonia: excursion guide* (Puura, V., Kalm, V., Puura, I., eds), Eesti Geoloogia Selts, Tallinn, p. 79–81.
- Männik, P., 2001. Distribution of conodonts. In: *Estonian Geological Sections. Bulletin, 3. Valga (10) drill core* (Pöldvere, A., ed.), Geological Survey of Estonia, Tallinn. p. 10–12.
- Männik, P., 2003. Distribution of conodonts. In: *Estonian Geological Sections. Bulletin, 5. Ruhnu (500) drill core* (Pöldvere, A., ed.), Geological Survey of Estonia, Tallinn, p. 17–23.
- Männil, R., Meidla, T., 1994. The Ordovician System of the East European Platform (Estonia, Latvia, Lithuania, Byelorussia, parts of Russia, the Ukraine and Moldova). In: *The Ordovician System of the East European Platform and Tuva (Southeastern Russia): Correlation Charts and Explanatory Notes* (Webby, B.D.R., Ross, R.J., Zhen, Y.Y., eds), IUGS Publication, **28**, A, p. 1–52.
- Männil, R., Rõõmusoks, A., 1984. The revision of the lithostratigraphic subdivision of the Ordovician of North Estonia. In: *Stratigraphy of the Lower Palaeozoic rocks in the East Baltic* (Männil, R., Mens, K., eds), Institute of Geology, Academy of Sciences of the Estonian SSR, Tallinn, p. 52–62.
- Männil, R., 1966. *Evolution of the Baltic Basin during the Ordovician*. Valgus, Tallinn, 200 pp. [in Russian with English summary]
- Männil, R., 1990. The Ordovician of Estonia. In: *Field Meeting Estonia 1990. An Excursion Guidebook* (Kaljo, D., Nestor, H., eds), Estonian Academy of Sciences, Tallinn, p. 11–20.
- Männil, R., Põlma, L., Hints, L., 1968. Stratigraphy of the Viruan and Harjuan deposits (Ordovician) in the central East Baltic. In: *Stratigraphy of the Lower Palaeozoic and Correlation with Other Regions* (Grigelis, A., ed.), Mintis, Vilnius, p. 81–110.
- Marshall, J. D., Brenchley, P. J., Mason, P., Wolff, G. A., Astini, R. A., Hints, L., Meidla, T., 1997. Global carbon isotopic events associated with mass extinction and glaciation in the late Ordovician. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **132**, 195–210.
- Meidla, T. 1996. Late Ordovician Ostracodes of Estonia. *Fossilia Baltica*, **2**, 1–222.
- Meidla, T., 1997. Hunneberg Stage. Billingen Stage. Volkhov Stage. Kunda Stage. In: *Geology and Mineral Resources of Estonia* (Raukas, A., Teedumäe, A., eds), Estonian Academy Publishers, Tallinn, p. 58–65.
- Meidla, T., Ainsaar, L., Hints, O., 2014. The Ordovician System in Estonia. In: *4th Annual Meeting of IGCP 591, Estonia, 10–19 June 2014. Abstracts and Field Guide* (Bauert, H., Hints, O., Meidla, T., Männik, P., eds). Tartu, University of Tartu, p. 116–122.
- Meidla, T., Ainsaar, L., Hints, O., 2023. About the Ordovician System in Estonia. In: *ISOS-14 Field Guide: The Ordovician of Estonia* (Hints, O., Toom, U., eds), TalTech Department of Geology, Tallinn, p. 9–14.
- Meidla, T., Truuver, K., Tinn, O., Ainsaar, L., 2020. Ostracods of the Ordovician–Silurian boundary beds: Jürjala core (Latvia) and its implications for Baltic stratigraphy. *Estonian Journal of Earth Sciences*, **69**, 233–247.
- Meidla, T., Hints, O., Ainsaar, L., 2023. Searching for the Ordovician–Silurian boundary in Estonia, Latvia and Lithuania. *Estonian Journal of Earth Sciences*, **72**, 70–73.
- Mickwitz, A., 1896. Über die Brachiopodengattung *Obolus* Eichwald. *Mém. Acad. Sci. Pétersbourg, sér. VIII, classe physico-mathématique*, **4**, 2, 216 pp.
- Miidel, A., Vaher, R., 1997. Neotectonics and recent crustal movements. In: *Geology and Mineral Resources of Estonia* (Raukas, A., Teedumäe, A., eds), Estonian Academy Publishers, Tallinn, p. 177–180.
- Miidel, A., 1997. Escarpments and waterfalls. In: *Geology and Mineral Resources of Estonia* (Raukas, A., Teedumäe, A., eds), Estonian Academy Publishers, Tallinn, p. 391–395.
- Nestor, H., Einasto, R., 1997. Ordovician and Silurian carbonate sedimentation basin. In: *Geology and Mineral Resources of Estonia* (Raukas, A., Teedumäe, A., eds), Estonian Academy Publishers, Tallinn, p. 192–204.
- Nõlvak, J., 1999. Ordovician chitinozoan biozonation of Baltoscandia. *Acta Universitatis Carolinae, Geologica*, **43**, 287–290.
- Nõlvak, J., Grahn, Y., 1993. Ordovician chitinozoan zones from Baltoscandia. *Review of Palaeobotany and Palynology*, **79**, 245–269.
- Nõlvak, J., 1986. *Rasprostranenie khitinozoi v razrezakh verkhnego ordovika Estonii*. AN ESSR Institut geologii, Tallinn. VINITI, N. 8526–B86.
- Nõlvak, J., Hints, O., Männik, P., 2006. Ordovician timescale in Estonia: recent developments. *Proceedings of the Estonian Academy of Sciences. Geology*, **55**, 95–108.
- Nõlvak, N., 1997. Ordovician. Introduction. In: *Geology and Mineral Resources of Estonia* (Raukas, A., Teedumäe, A., eds), Estonian Academy Publishers, Tallinn, p. 52–55.
- Öpik, A., 1930. Brachiopoda Protremata der estländischen ordovizischen Kukruse-Stufe. *Tartu Ülikooli Geoloogia Instituudi Toimetised*, **20**, 261 pp.
- Öpik, A., 1934. Über Klitamboniten. *Acta et commentationes Universitatis Tartuensis*, **A 26**, 1190.
- Orviku, K., 1940. Lithologie der Tallinna-serie (Ordovizium, Estland). *Acta et commentationes Universitatis Tartuensis*, **A 36**, 1–216.
- Orviku, K., 1960. On lithostratigraphy of the Toila and Kunda stages in Estonia. *Geoloogia Instituudi Uurimused*, **5**. Eesti Teaduste Akadeemia, Tallinn. p. 45–87.
- Paiste, T., Männik, P., Meidla, T., 2022. Sandbian (Late Ordovician) conodonts in Estonia: distribution and biostratigraphy. *GFF*, **69**, 76–90.
- Paiste, T., Männik, P., Meidla, T., 2023. Emended Sandbian (Ordovician) conodont biostratigraphy in Baltoscandia and a new species of *Amorphognathus*. *Geological Magazine*, **160**, 411–427.
- Põlma, L., 1982. *Comparative Lithology of the Ordovician Carbonate Rocks in the Northern and Middle East Baltic*. Valgus, Tallinn. 164 pp.
- Puura, I., Viira, V., 1999. Chronostratigraphy of the Cambrian–Ordovician boundary beds in Baltoscandia. *Acta Universitatis Carolinae Geologica*, **43**, 5–8.
- Raukas, A., Teedumäe, A. (eds), 1997. *Geology and Mineral Resources of Estonia*. Estonian Academy Publishers, Tallinn. 436 pp.
- Raymond, P. E., 1916. The correlation of the Ordovician strata of the Baltic basin with those of North America. In: Raymond, P. E., Twenhofel, W. (eds), Expedition to the Baltic provinces of Russia and Scandinavia, 1914. *Bulletin of the Museum of Comparative Zoology at Harvard College*, **56**, 179–286.
- Remane, J., 2003. Chronostratigraphic correlations: their importance for the definition of geochronologic units. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **196**, 7–18.
- Resheniya..., 1965. *Resolution of the joint discussion on establishing an unitary Precambrian and Palaeozoic scheme for the Russian Platform*, 1962. VSEGEI, Leningrad, 79 pp.
- Resheniya..., 1978. *Resolution of inter-regional stratigraphic discussion on establishing an unitary stratigraphic scheme for the East Baltic region*, 1976. Lithuanian NIGRI, Leningrad, 86 pp.
- Resheniya..., 1987. *Resolution of the joint stratigraphic discussion on the Ordovician and Silurian of the East European Platform*, 1984. VSEGEI, Leningrad, 114 pp.
- Rong, J.-y., Chen, X., Harper, D. A. T., 2002. The latest Ordovician Hirnantia fauna (Brachiopoda) in time and space. *Lethaia*, **35**, 231–249.
- Rõõmusoks, A., 1983. *Eesti aluspõhja geoloogia*. Valgus, Tallinn. 224 pp.
- Rõõmusoks, A., Puura, V., Raukas, A., Mark-Kurik, E., 1997. History of geological research. In: *Geology and Mineral Resources of Estonia* (Raukas, A., Teedumäe, A., eds), Estonian Academy Publishers, Tallinn, p. 15–26.
- Schmidt, F., 1858. Untersuchungen über die Silurische Formation von Ehstland, Nord-Livland und Oesel. *Archiv für die Na-*

- turkunde Liv- Ehst- and Kurlands, ser. 1, 2*, Dorpat, 248 pp.
- Schmidt, F., 1881. Revision der ostbaltischen silurischen Tri-lobiten nebst geognostischer Übersicht des ostbaltischen Silurgebiets. Abt. 1. *Mémoires de l'Académie Impériale des Sciences de St. Petersburg, ser. 7, 30, 1*, 238 pp.
- Schmidt, F., 1897. *Excursion durch Estland. Guide des excursions du VII Congrès Géologique International, St. Pétersbourg*, 1897, 21 pp.
- Strangways, W. T. H. F., 1821. Geological sketch of the environs of Petersburg. *Transactions of the Geological Society of London, 5*, 392–458.
- Suuroja, K. 2006. *Baltic Klint in North Estonia. A Symbol of Estonian Nature*. Ministry of Environment, 224 pp.
- Tinn, O., Meidla, T., 2001. Middle Ordovician ostracods from the Lanna and Holen Limestones, south-central Sweden. *GFF, 123*, 129–136.
- Tuuling, I., 1998. Shipborne geophysical study of an Ordovician–Silurian carbonate platform, Faro Hiiumaa area, northeastern Baltic Sea. *Meddelanden från Stockholms Universitets Institution för Geologi och Geokemi, 301*, 21 pp.
- Tuuling, I., Flodén, T., 2013. Silurian reefs off Saaremaa and their extension towards Gotland, central Baltic Sea. *Geological Magazine, 150*, 923–936.
- Wahl, A. von, 1923. Mitteilungen über die Geologie von Borkholm und seiner Umgebung. *Loodusuurijate Seltsi aruan-ded, 29*, 23–29.
- Webby, B. D., 1998. Steps towards a global standard for Ordovician stratigraphy. *Newsletters on Stratigraphy, 36*, 1–33.

XI Baltic Stratigraphical Conference. Abstracts and Field Guide

Edited by Olle Hints, Peep Männik and Ursula Toom

XI Baltic Stratigraphical Conference, Tartu and Arbavere, Estonia (August 19–21, 2024)

Post-conference Field Excursion (August 22–25, 2024)


The conference and field excursion are organised by:

Geological Society of Estonia

Tallinn University of Technology, Department of Geology

University of Tartu, Department of Geology

Geological Survey of Estonia



XI Baltic Stratigraphical Conference is a regional meeting of the International Geoscience Programme Project 735: "Rocks and the Rise of Ordovician Life"

© 2024 Authors. This publication is distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0>), if not indicated otherwise.

Electronic copy available at: <https://geologia.info/reference/49701>

Published by the Geological Society of Estonia

Printed by Alfapress OÜ

ISBN 978-9949-9957-5-2 (print)

ISBN 978-9949-9957-6-9 (pdf)

Tallinn, 2024