

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/272437661>

Bryozoans from the Lower Silurian (Wenlockian) Steinsfjorden Formation of Ringerike, southern Norway

Article in *Bulletin of Geosciences* · January 2015

DOI: 10.3140/bull.geosci.1521

CITATIONS

2

READS

436

2 authors:



Andrej Ernst

University of Hamburg

212 PUBLICATIONS 1,781 CITATIONS

SEE PROFILE



Hans Arne Nakrem


University of Oslo

115 PUBLICATIONS 1,535 CITATIONS

SEE PROFILE

Bryozoans from the lower Silurian (Wenlock) Steinsfjorden Formation of Ringerike, southern Norway

ANDREJ ERNST & HANS ARNE NAKREM



A stenolaemate bryozoan fauna from the lower Silurian (Wenlock) Steinsfjorden Formation of the Ringerike district, southern Norway contains 10 species (2 cystoporates, 6 trepostomes, and 2 cryptostomes). Six species are new: *Heterotrypa ringerikensis* sp. nov., *Trematopora maculata* sp. nov., *Amplexopora crassiparietum* sp. nov., *A. evae* sp. nov., *Orthopora worsleyi* sp. nov., and *Mediaporina kiaeri* sp. nov. One species is described in open nomenclature: *Eridotrypella* sp. Trepostome bryozoans dominate this assemblage, both by diversity and abundance. The bryozoans occur in carbonates associated with evaporite minerals and sedimentary structures indicating very shallow depositional settings. • Key words: lower Silurian, Bryozoa, Norway, taxonomy.

ERNST, A. & NAKREM, H.A. 2015. Bryozoans from the lower Silurian (Wenlock) Steinsfjorden Formation of Ringerike, southern Norway. *Bulletin of Geosciences* 90(1), 65–87 (13 figures, 10 tables). Czech Geological Survey, Prague. ISSN 1214-1119. Manuscript received June 5, 2014; accepted in revised form September 11, 2014; published online December 3, 2014; issued January 26, 2015.

Andrej Ernst, Institut für Geologie, Universität Hamburg, Bundesstr. 55, D-20146 Hamburg, Germany; Andrej.Ernst@uni-hamburg.de • Hans Arne Nakrem, Natural History Museum (Geology), University of Oslo, P.O. Box 1172 Blindern, NO-0318 Oslo, Norway; h.a.nakrem@nhm.uio.no

The purpose of this study is to describe the bryozoan fauna of the lower Wenlock Brattstad Member, Steinsfjorden Formation of the Ringerike district, Oslo Region, Norway (Fig. 1). The bryozoans present in these beds have been known since the work by Kiær (1908), but they have not until now been given a systematic treatment. The number of previously known species is low, and their distribution in Ringerike can add very little to the already established biostratigraphy of the area.

The Brattstad Member (upper “Etage” 9e and 9f of Kiær 1908) is 30 m thick in the studied areas of Nes bus stop, Ødegårdsviken and Åsa (Fig. 2). The lithology consists of medium to thickly bedded limestones with interbedded marls and dolomitic limestones. The limestones contain a relatively rich and varied fauna with thick massive favositid and oncolitic biostromes (the “*Favosites* Limestone” of Kiær 1908) near the top of the member from which the bryozoan material derives. Kiær (1908, p. 113) lists fossils from these beds (“Etage 9f”) in various localities in the Ringerike district, but only one bryozoan taxon, “*Monticuliporen* [*Amplexopora*(?) sp.]”, is mentioned. Whitaker (1977, pp. 24–26) also refers to the bryozoan-rich beds as “polyzoan (*Amplexopora*)” beds. Spjeldnæs (1982) describes the bryozoan occurrences in these beds as “colonies ranging from only a few cm in diameter to small bioherms with three-dimensional frameworks”. Bryozoans are even found in the lowermost levels of the overlying red

beds of the Ringerike Group (late Wenlock to Ludlow age). The age of the bryozoan beds are Sheinwoodian to Homerian as based on occurrences of ostracodes and brachiopods as well as agnathans and thelodonts (on directly overlying beds) (Worsley *et al.* 1983, 2011). Geochemical analyses of interbedded bentonites (Hetherington *et al.* 2011) support these datings.

Depositional environment

Alternations of marls, red argillaceous dolomites, dolomitic limestones, and limestones, with occurrence of oospirite, oncolites, algal laminated dolomitic limestones with mud cracks, stromatolites as well as evaporitic minerals (Fig. 3B) indicate developments of shallow subtidal to supratidal environments (Olaussen 1981). The fossil components are characterized by many individuals belonging to few taxa, notably gastropods, ostracods, brachiopods and rare trilobites (Kiær 1908, Olaussen 1981). Tabulate corals (Fig. 3C) and echinoderms are locally present.

The common microfacies are represented by floatstones with abundant bryozoans (mainly trepostomes (Fig. 3A, F), locally poorly washed rudstones (Fig. 3E), and bryozoan and algal bindstones (Fig. 3B–D). The matrix is usually micrite-rich, containing various skeletal grains (Fig. 3G). Calcimicrobes are common on some localities,




Figure 1. Map of the Ringerike District showing sample localities and major structural features. (A) – Åsa locality, (B) – Ødegårdsviken locality, (C) – Nes bus stop locality.

Per.	Epoch/Age		Baltic sta.	Ringerike lithostrat.
Silurian	Ludlow	Corstian	Paadla	Ringerike Group
	Wenlock	Homerian	Rootsiküla	Steinsfjorden Fm. Ranberget Mbr.
			Jaagarahu	Brattstad Mbr. Sjøvoll Mbr.
		Sheinwoodian	Jaani	Braksøya Fm.
			Llandov.	Telychian

Figure 2. Wenlock stratigraphy of the Ringerike District (based on Melchin *et al.* 2012; Worsley *et al.* 1983, 2011).

represented by various *Girvanella* and *Rothpletzella* species. Microbial crusts are common, often forming oncoids (Fig. 3C).

The faunal composition and microfacies characteristics are reduced due to stressed conditions in an environment with fluctuating salinities.

Material and methods


Bryozoan samples were collected from the localities Nes bus stop, Ødegårdsviken and Åsa (see Fig. 1). The investigated bryozoans were studied from thin sections using a transmitted light binocular microscope. 107 oriented and non-oriented thin sections were used. The material is housed at the Natural History Museum (Geology), Oslo, under numbers PMO 221.750–221.841.

Morphological character terminology is adopted from

Anstey & Perry (1970) for trepostomes, and Hageman (1993) for cryptostomes. The following morphologic characters were measured for statistical use (Fig. 4): Branch Width, Branch Thickness, Exo- (Endo-) zone Width, Autozoecial Aperture Width, Autozoecial Aperture Spacing (Along Branch/diagonally for cryptostomes), Acanthostyle Diameter, Lunarium Width/Length/Thickness, Vesicle Diameter/Spacing/Number per Aperture, Wall Thickness in Exozone, and Macular Diameter (Spacing), Autozoecial Diaphragm Spacing, Meso- (Exila-, Hetero-) zooecia Width, Mesozoecial Diaphragm Spacing.

The spacing of structures was measured as the distance between centres. Additional quantitative characters include the Number of Mesozoecia, Exilazoecia and Acanthostyles surrounding each autozoecial aperture. Statistics were summarized using arithmetic mean, sample standard deviation, coefficient of variation, and minimum and maximum values.

Figure 3. Carbonate microfacies of the Steinsfjorden Formation of Ringerike, southern Norway. • A – bryozoan floatstone, Nes bus stop, PMO 221.766. • B – bryozoan bindstone (*Trematopora maculata*, *Fistulipora nummulina*). Arrow – relicts of gypsum crystals, Ødegårdsviken, PMO 221.820. • C – coral-oncolitic floatstone, Åsa, PMO 221.813. • D – bryozoan bindstone (*Trematopora maculata*, *Fistulipora nummulina*), Ødegårdsviken, PMO 221.824. • E – poorly washed rudstone, Ødegårdsviken, PMO 221.779. • F – bryozoan floatstone, Ødegårdsviken, PMO 221.792. • G – matrix, Nes bus stop, PMO 221.766.



Results and discussion

The bryozoan colonies are generally well preserved and show little signs of abrasion. Large, thin branches up to 5 mm in diameter are often not broken, the whole fauna has only been transported a short distance, and they may represent time intervals of more normal marine conditions. However, they are not considered preserved *in situ*. The bryozoans are therefore interpreted as typical for the depositional environment in which the sediments were formed. The three bryozoans species previously known have a Wenlock or Ludlow stratigraphic distribution elsewhere. The new species described herein have many similarities with species known from time equivalent strata in Europe and the USA, but only further research may reveal their biostratigraphic as well as biogeographic value.

Systematic palaeontology

Phylum Bryozoa Ehrenberg, 1831
 Class Stenolaemata Borg, 1926
 Order Cystoporata Astrova, 1964
 Family Fistuliporidae Ulrich, 1882

Genus *Fistulipora* M'Coy, 1849

Type species. – *Fistulipora minor* M'Coy, 1849. Carboniferous; England.

Diagnosis. – Massive, encrusting or ramose colonies. Cylindrical autozooecea with thin walls and complete diaphragms. Apertures rounded, possessing horse-shoe shaped lunaria. Autozooecea separated by the extrazooecial vesicular skeleton.

Comparison. – *Fistulipora* M'Coy, 1849 differs from *Eridopora* Ulrich, 1882 in having rounded, horseshoe-shaped lunaria instead of triangular ones. Furthermore, *Eridopora* develops persistently encrusting colonies, whereas *Fistulipora* may also develop massive and branched colonies.

Occurrence. – Ordovician to Permian; worldwide.

Fistulipora nummulina Nicholson & Foord, 1885

Figure 5A–E, Table 1

- 1885 *Fistulipora nummulina* Nicholson & Foord, pp. 506–507, text-fig. 4, pl. 15, figs 2–2b.
- 1885 *Fistulipora dobunica* Nicholson & Foord, p. 511, pl. 17, figs 3–3b.
- 1969 *Fistulipora nummulina* Nicholson & Foord, 1885. – Owen, pp. 627–628, pl. 114, figs 1, 2.

Table 1. Measurements of *Fistulipora nummulina* Nicholson & Foord, 1885. Abbreviations: N – number of measurements; X – mean; SD – standard deviation; CV – coefficient of variation; MIN – minimal value; MAX – maximal value.

	N	X	SD	CV	MIN	MAX
Aperture width, mm	33	0.16	0.027	16.46	0.12	0.22
Aperture spacing, mm	31	0.24	0.035	14.41	0.17	0.32
Vesicle width, mm	23	0.10	0.031	32.55	0.03	0.15
Vesicle spacing, mm	25	0.10	0.028	28.98	0.06	0.15
Vesicles per aperture	8	6.0	1.414	23.57	4.0	8.0

Material. – PMO 221.756, PMO 221.758, PMO 221.790, PMO 221.764, PMO 221.817, PMO 221.820, PMO 221.822, PMO 221.824, PMO 221.827, PMO 221.829.

Description. – Encrusting colonies, 0.25–0.85 mm in thickness. Autozooecea growing from 0.002–0.008 m thick epitheca, bending sharply at their bases towards colony surface. Autozooeccial apertures circular to oval. Basal diaphragms rare to absent. Lunaria well developed, horseshoe shaped to slightly triangular, 0.06–0.08 mm wide, 0.03–0.04 mm long and 0.03–0.04 mm thick. Vesicles abundant, 4–8 surrounding each autozooeccial aperture, irregularly shaped, medium in size, often large at the base, polygonal in tangential section, box-like to hemispheric, with plane or concave roofs in exozone. Autozooeccial walls granular, 0.003–0.005 mm thick. Extensive extrazooecial material in exozone developed. Maculae not observed.

Remarks. – *Fistulipora nummulina* Nicholson & Foord, 1885 is similar to *F. parva* Hennig, 1908 from the Wenlock of Gotland, but differs by smaller autozooeccial apertures (aperture width 0.12–0.22 mm vs 0.22–0.25 mm in *F. parva*). *Fistulipora nummulina* differs from *F. ternavensis* Astrova, 1965 from the lower Silurian (Wenlock) of Moldova in smaller and closer spaced autozooeccial apertures (aperture width 0.12–0.22 mm vs 0.18–0.23 mm in *F. ternavensis*).

Occurrence. – Much Wenlock Limestone Formation, Silurian, Wenlock, Homerian; Dudley, England. Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian; Nes bus stop, Ringerike, Norway.

Fistulipora crustula Bassler, 1906

Figures 5F–J, 6A, B, Table 2

- 1906 *Fistulipora crustula* Bassler, p. 24, pl. 7, figs 7–10, pl. 8, figs 16, 17, pl. 23, fig. 15.

Material. – PMO 221.775–221.776.




Figure 4. Some important measurements made on the trepostome and cystoporate bryozoans. Abbreviations: ADB – aperture spacing, AW – autozoecial aperture width, LuL – lunarium length, LuW – lunarium width, LuT – lunarium thickness, VSp – vesicle spacing, DSp – autozoecial diaphragm spacing, MD – meso- (exila-, hetero-) zooecial width, MDSp – mesozooecial diaphragms spacing, AcD – acanthostyles diameter.

Description. – Encrusting colonies, 0.65–1.38 mm in thickness. Autozoecia growing from 0.015–0.030 mm thick epitheca, bending sharply at their bases towards colony surface. Autozoecial apertures circular to oval. Basal diaphragms rare to common, thin, deflected proximally. Lunaria well developed, horseshoe shaped. Vesicles irregularly shaped, medium in size, polygonal in tangential section, box-like to hemispheric, with concave roofs in exozone, 5–11 surrounding each autozoecial aperture. Autozoecial walls granular, 0.003–0.005 mm thick. Maculae consisting of vesicular skeleton common, 0.30–0.40 mm in diameter. Macrozoecia locally occurring.

Remarks. – *Fistulipora crustula* Bassler, 1906 is similar to *F. crustuliformis* Astrova, 1959 from the upper Silurian (Ludlow) of Tuva (Russia), but differs from the latter in smaller autozoecia (aperture width 0.18–0.25 mm vs 0.22–0.26 mm in *F. crustuliformis*).

Occurrence. – Rochester Shale, Wenlock, lower Silurian; USA. Steinsfjorden Formation, Brattstad Member, Silur-

ian, Wenlock, Sheinwoodian–Homerian; Ødegårdsviken, Ringerike, Norway.

Order Trepostomata Ulrich, 1882
Suborder Halloporina Astrova, 1965
Family Heterotrypidae Ulrich, 1890

Genus *Heterotrypa* Nicholson, 1879

Type species. – *Monticulipora frondosa* d’Orbigny, 1850. Upper Ordovician (Cincinnatian); North America.

Diagnosis. – Frondose, ramose or less commonly encrusting colonies. Autozoecial walls variably thick. Zoecial boundaries form a conspicuous dark line in inner exozones and in a broad zone of abutting laminae or are completely obscured in outer exozones. Walls generally amalgamated. Diaphragms are generally few in endozones, but are moderately abundant in some species. In exozones, autozoecial diaphragms are closely and regularly spaced, thin, planar and perpendicular to

Table 2. Measurements of *Fistulipora crustula* Bassler, 1906. Abbreviations as in Table 1.

	N	X	SD	CV	MIN	MAX
Aperture width, mm	25	0.21	0.019	8.95	0.18	0.25
Aperture spacing, mm	25	0.25	0.043	17.02	0.18	0.35
Vesicle width, mm	25	0.07	0.023	31.10	0.03	0.13
Vesicle spacing, mm	25	0.08	0.020	24.97	0.04	0.11
Vesicles per aperture	6	8.2	2.041	24.99	5.0	11.0
Lunarium width, mm	7	0.15	0.035	23.41	0.11	0.20
Lunarium length, mm	7	0.035	0.012	33.35	0.025	0.050
Lunarium thickness, mm	7	0.033	0.014	43.25	0.015	0.050
Macrozoecia aperture width, mm	10	0.30	0.019	6.41	0.28	0.33

the zooecial walls. Intermonticular mesozoecia range from abundant and regularly arranged to scattered or absent. Mesozoecia commonly develop moniliform chambers at proximal ends and tend to become smaller or are terminated distally within exozones. Diaphragms in mesozoecia noticeably thicker and more closely spaced than autozoecial diaphragms. Acanthostyles common, small to moderately large, occurring both in endozone and exozone. Monticules generally have a central cluster of a few mesozoecia.

Remarks. – *Heterotrypa* differs from the most similar genus *Dekayia* Milne-Edwards & Haime, 1851 by having fewer and smaller acanthostyles as well as more abundant mesozoecia.

Occurrence. – Ordovician to Devonian; worldwide.

***Heterotrypa ringerikense* sp. nov.**

Figures 6C–F, 7A–E, Table 3

Etymology. – The species has its name after Ringerike, an area SW of the Oslo Region, which is famous for its rich Silurian bryozoan-bearing rocks.

Holotype. – PMO 202.721B.

Paratypes. – PMO 221.763, PMO 221.773–221.777, PMO 221.785–221.786, PMO 221.799, PMO 202.721(A, C–E).

Type locality. – Ødegårdsviken, Ringerike, Norway.

Type stratum. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian.

Diagnosis. – Ramose branched colonies; secondary overgrowth common; autozoecial diaphragms rare to common, thin; mesozoecia rare to common; acanthostyles moderately large, 2–7 surrounding each autozoecial aperture; macroacanthostyles present; maculae consisting of macrozoecia.

Description. – Ramose branched colonies, branch diameter 2.4–5.0 mm. Exozone distinct, 0.45–1.25 mm wide, endozone 1.30–2.88 mm wide. Secondary overgrowth common, 0.65–1.00 mm in thickness. Autozoecia long, polygonal in cross section in endozone, bending sharply in exozone. Autozoecial apertures polygonal. Autozoecial diaphragms rare to common, thin, concentrated mainly in the transition between exo- and endozones. Mesozoecia rare to common, originating at base of exozone, beaded in places of development of diaphragms. Diaphragms in mesozoecia straight. Acanthostyles moderately large, prominent, having distinct hyaline cores, 2–7 surrounding each autozoecial aperture. Macroacanthostyles locally present. Autozoecial walls 0.005–0.010 mm thick, granular-prismatic in endozone; showing reversal V-shaped lamination, integrated with locally visible serrated dark border between zooecia, locally weakly monilae-shaped thickened, 0.013–0.050 mm thick in exozone. Maculae consisting of macrozoecia.

Remarks. – *Heterotrypa ringerikense* sp. nov. differs from *H. enormis* Astrova, 1959 from the upper Silurian (Ludlow) of western Tuva (Russia) by larger and more abundant acanthostyles (acanthostyle diameter 0.025–0.065 mm vs 0.03 mm in *H. enormis*; 2–7 acanthostyles around each aperture vs 3–4 in *H. enormis*). *Heterotrypa ringerikense* sp. nov. differs from *H. ovata* Astrova, 1957 from the upper Silurian (Ludlow) of western Tuva (Russia) by less abundant mesozoecia and by larger and more abundant acanthostyles (2–7 acanthostyles around each aperture vs 1–2 in *H. ovata*).

Occurrence. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian; Ødegårdsviken, Ringerike, Norway.

Genus *Asperopora* Owen, 1969

Type species. – *Callopora aspera* Hall, 1852. Silurian (Wenlock); USA (New York), England.

Figure 5. A–E – *Fistulipora nummulina* Nicholson & Foord, 1885. • A, B – encrusting colony, PMO 221.764. • C – longitudinal section, PMO 221.820. • D – tangential section, PMO 221.817. • E – tangential section, PMO 221.764. • F–J – *Fistulipora crustula* Bassler, 1906. • F, G – longitudinal section, PMO 221.775. • H – longitudinal section, PMO 221.776. • I, J – tangential section, PMO 221.775.




Table 3. Measurements of *Heterotrypa ringerikense* sp. nov. Abbreviations as in Table 1.

	N	X	SD	CV	MIN	MAX
Branch width, mm	6	3.50	1.172	33.51	2.40	5.00
Exozone width, mm	6	0.74	0.312	42.08	0.45	1.25
Endozone width, mm	6	2.01	0.612	30.40	1.30	2.88
Aperture width, mm	55	0.17	0.025	15.13	0.11	0.22
Aperture spacing, mm	55	0.20	0.031	14.94	0.15	0.29
Aperture width, mm (macular)	20	0.24	0.025	10.47	0.2	0.28
Aperture spacing, mm (macular)	20	0.30	0.044	14.63	0.24	0.42
Mesozoecia width, mm	50	0.06	0.021	37.98	0.03	0.10
Acanthostyle diameter, mm	45	0.044	0.009	20.22	0.025	0.065
Macroacanthostyle diameter, mm	20	0.067	0.012	17.22	0.043	0.088
Exozonal wall thickness, mm	45	0.030	0.010	31.96	0.013	0.050
Acanthostyles per aperture	45	3.7	1.424	38.38	2.0	7.0
Mesozoecial diaphragms spacing, mm	10	0.10	0.025	25.40	0.05	0.14

Diagnosis. – Colonies incrusting, massive or branched ramose; autozoecia tubular; autozoecial apertures rounded to petaloid due to inflecting acanthostyles; autozoecial diaphragms few to many; mesozoecia abundant, originating in exozones, sometimes cystose, containing thin diaphragms; acanthostyles commonly large and abundant, with clear cores, inflecting autozoecial apertures; autozoecial walls moderate in thickness, finely laminated, merged, irregularly thickened in exozone (modified after Owen 1969).

Remarks. – *Asperopora* Owen, 1969 is similar to *Leioclema* Ulrich, 1882 but differs by irregularly thickened walls in exozone. In general, *Asperopora* has thinner walls and smaller acanthostyles. *Asperopora* differs from *Trematopora* Hall, 1852 by having mesozoecia not sealed by calcitic material. Moreover, *Trematopora* develops exceptionally ramose-branched colonies, whereas *Asperopora* forms also encrusting and massive ones.

Occurrence. – Upper Ordovician to upper Silurian; Europe, USA.

***Asperopora bellum* (Pushkin, 1976)**

Figure 7F–I, Table 4

1976 *Lioclema bellum* Pushkin, pp. 30–31, text-fig. 13, pl. 11, fig. 3.

Material. – PMO 221786–221.787, PMO 221.793, PMO 221.798, PMO 221.800.

Description. – Encrusting colonies, 0.7–1.0 mm thick. Autozoecia budding from a thin epitheca, growing a short distance parallel to the substrate, then bending sharply to the colony surface. Autozoecial apertures rounded-polygonal. Autozoecial diaphragms common, straight, thin. Acanthostyles abundant, 1–3 surrounding each aperture, originating at the base of exozone, having distinct calcite cores and dark, laminated sheaths, deeply indenting into autozoecial space. Mesozoecia abundant, 6–11 surrounding each autozoecial aperture, rounded-polygonal in transverse section, originating at the base of exozone. Autozoecial walls laminated, merged, 0.020–0.050 mm thick. Maculae not observed.

Remarks. – *Asperopora bellum* (Pushkin, 1976) differs from *A. asperum* (Hall, 1852) in larger autozoecial apertures (average aperture width 0.14 mm vs 0.12 mm in *A. asperum*), and in less abundant mesozoecia (6–11 mesozoecia per aperture vs 9–14 in *A. asperum*). *Asperopora bellum* differs from *A. multipora* (Bassler, 1906) from the Rochester Shale of USA in smaller autozoecial apertures (aperture width 0.10–0.18 mm vs 0.15–0.20 mm in *A. multipora*), and fewer acanthostyles (1–3 vs 4–7 per aperture in *A. multipora*).

Occurrence. – Ludlow, Silurian; Brest Depression, Belarus. Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian; Ødegårdsviken, Ringerike, Norway.

Family Trematoporidae Miller, 1889

Genus *Trematopora* Hall, 1852

Type species. – *Trematopora tuberculosa* Hall, 1852; lower Silurian (Niagaran); North America.

Diagnosis. – Ramose colonies, often originating from encrusting base. Autozoecial apertures oval to rounded with peristomes. Diaphragms usually rare, often absent in endozone. Abundant mesozoecia with abundant diaphragms, thin-walled and beaded in initial parts of exozone, near colony surface becoming thick-walled. Mesozoecial apertures often completely covered by laminated skeleton. Acanthostyles abundant. Walls thin in endozone, thickened in

Figure 6. A, B – *Fistulipora crustula* Bassler, 1906, tangential section, PMO 221.775. • C–F – *Heterotrypa ringerikensis* sp. nov.; C, D – longitudinal section, holotype PMO 202.721B; E – transverse section, holotype PMO 202.721B; F, G – longitudinal section of encrusting colony, paratype PMO 202.721A.




Table 4. Measurements of *Asperopora bellum* (Pushkin, 1976). Abbreviations as in Table 1.

	N	X	SD	CV	MIN	MAX
Aperture width, mm	36	0.14	0.023	16.74	0.10	0.18
Aperture spacing, mm	36	0.20	0.031	15.50	0.14	0.26
Mesozooecea width, mm	36	0.05	0.015	32.18	0.02	0.07
Mesozooecea per aperture	30	7.4	1.329	17.96	6.0	11.0
Acanthostyle diameter, mm	36	0.037	0.007	17.90	0.025	0.060
Acanthostyles per aperture	30	2.3	0.596	25.91	1.0	3.0
Mesozooeccial diaphragms spacing, mm	21	0.09	0.029	30.43	0.06	0.17

Table 5. Measurements of *Trematopora maculata* sp. nov. Abbreviations as in Table 1.

	N	X	SD	CV	MIN	MAX
Branch width, mm	10	3.7	0.685	18.61	2.7	4.6
Exozone width, mm	10	0.8	0.198	25.28	0.4	1.1
Endozone width, mm	10	2.1	0.364	17.21	1.4	2.6
Aperture width, mm	79	0.16	0.025	15.43	0.11	0.22
Aperture spacing, mm	79	0.22	0.036	16.73	0.15	0.36
Aperture width, mm (macular)	15	0.25	0.025	9.74	0.23	0.33
Aperture spacing, mm (macular)	15	0.33	0.058	17.69	0.27	0.52
Mesozooecea width, mm	70	0.065	0.019	28.76	0.033	0.120
Mesozooecea per aperture	30	2.9	0.960	33.09	1.0	4.0
Acanthostyle diameter, mm	70	0.052	0.010	19.77	0.030	0.088
Acanthostyles per aperture	70	2.7	0.823	30.47	1.0	4.0
Mesozooeccial diaphragms spacing, mm	70	0.10	0.024	23.39	0.06	0.19
Autozooeccial diaphragms spacing, mm	30	0.16	0.063	38.50	0.10	0.30
Exozonal wall thickness, mm	50	0.037	0.010	27.07	0.018	0.058

peripheral parts of exozone displaying obliquely laminated microstructure.

Remarks. – *Trematopora* Hall, 1852 differs from *Batosotoma* Ulrich, 1882 by having oval to rounded autozooeccial apertures and abundant mesozooecea covered with skeletal material, from *Eridotrypa* Ulrich, 1893 by having autozooecea that bend sharply in exozone, possess rounded apertures and are arranged irregularly on the colony surface, as well as by abundant acanthostyles.

Occurrence. – Ordovician to Silurian, worldwide.

***Trematopora maculata* sp. nov.**

Figure 8A–H, Table 5

Etymology. – The species is named after presence of maculae of macrozooecea.

Holotype. – PMO 221.816.

Paratypes. – PMO 221.783–PMO 221.784, PMO 221.786–PMO 221.787, PMO 221.791, PMO 221.799, PMO 221.815, PMO 221.817–221.841.


Type locality. – Ødegårdsviken, Ringerike, Norway.

Type stratum. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian.

Diagnosis. – Branched colonies with distinct exozones; secondary overgrowth common; autozooeccial diaphragms rare to common; mesozooecea common to abundant in exozone, beaded, 2–4 surrounding each autozooeccial aperture; acanthostyles moderately large, 1–4 surrounding each autozooeccial aperture; autozooeccial walls laminated with locally visible zooeical border, locally monilae-shaped thickened, maculae of few macrozooecea.

Description. – Ramose branched colonies, branch diameter 2.7–4.6 mm. Exozone distinct, 0.4–1.1 mm wide, endozone 1.4–2.6 mm wide. Secondary overgrowth common. Autozooecea long, polygonal in cross section in endozone, bending sharply in exozone. Autozooeccial apertures rounded to slightly angular. Autozooeccial diaphragms rare to common, thin, concentrated mainly in the transition between exo- and endozones. Mesozooecea common to abundant, originating at base of exozone, beaded in places of development of diaphragms, 2–4 surrounding each autozooeccial aperture. Diaphragms in mesozooecea straight, abundant. Acanthostyles moderately large, prominent, having distinct hyaline cores, 1–4 surrounding each autozooeccial aperture. Autozooeccial walls 0.003–0.005 mm thick, granular-prismatic in endozone; showing reversal V-shaped lamination, integrated with locally visible dark border between zooecea, locally monilae-shaped thickened, 0.02–0.06 mm thick in exozone. Maculae formed by few macrozooecea.

Figure 7. A–E – *Heterotrypa ringerikensis* sp. nov.; A, B – tangential section, paratype PMO 202.721A.; C, D – branch transverse section, paratype PMO 221.763.; E – tangential section showing macroacanthostyles, paratype PMO 202.721E. • F–I – *Asperopora bellum* (Pushkin, 1976), PMO 221.793; F – tangential section; G, H – longitudinal section; I – tangential section. • J – *Trematopora maculata* sp. nov., longitudinal section, holotype PMO 221.816.



Remarks. – *Trematopora maculata* sp. nov. is similar to *T. beikhemensis* Astrova, 1959 from the Wenlock of western Tuva (Russia), but differs by presence of macrozoecia, broader exozone, and thicker walls. *Trematopora maculata* sp. nov. differs from *T. cristata* Kopaevich, 1984 from the Wenlock of Mongolia by presence of macrozoecia and less elongated apertures.

Occurrence. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian; Ødegårdsviken, Ringerike, Norway.

Suborder Amplexoporina Astrova, 1965

Family Amplexoporidae Miller, 1889

Genus *Amplexopora* Ulrich, 1882

Type species. – *Atactopora septosa* Ulrich, 1879. Upper Ordovician; USA.

Diagnosis. – Ramose, frondescent, incrusting or massive colonies. Monticules generally well developed. Wall structure commonly appearing integrate in the exozone, sometimes amalgamate. Laminae from adjacent zooecia intersect a sharply defined zooecial boundary at angles of less than 90° to form a V-shaped pattern pointing distally. Distinct zooecial linings present in several species. Diaphragms are extremely variable in thickness, curvature, parallelism and spacing, with compound and cystoidal diaphragms and cystiphagms common in many species. Mesozooecia lacking or few; early chambers are beaded and later diaphragms regularly and closely spaced. Acanthostyles usually abundant and of two types: those that are concentrated generally in zooecial and extend throughout the width of the exozone, occurring in all species; and additional acanthostyles that are concentrated between zooecial corners and extend through a part of the exozone width occurring in some species. These additional acanthostyles cause inflections of zooecial walls.

Remarks. – *Amplexopora* differs from *Monotrypa* by abundant diaphragms and wall microstructure with sharply defined zooecial boundary.

Boardman (1960) revised the genus *Amplexopora* and designated *Atactopora septosa* Ulrich, 1879 as the type species. According to the emended diagnosis of Boardman (1960) (followed here), *Amplexopora* possesses mesozooecia with abundant diaphragms. On the contrary, Astrova

(1978, p. 105) mentioned exilazooecia in *Amplexopora*, tube-like heterozooecia without or with few diaphragms. Based on this character she defined the suborder Amplexoporina, which comprises trepostome bryozoans with exilazooecia. Furthermore, Astrova (1978) re-designated *Amplexopora cingulata* Ulrich, 1882 as the type species of *Amplexopora*. According to Boardman (1960, p. 17) this species was never described or figured (except in Astrova 1978, pl. 17, fig. 2).

Occurrence. – Lower Ordovician–lower Silurian; North America, Australia, Siberia, Europe.

Amplexopora crassiparietum sp. nov.

Figure 9A–F, Table 6

Etymology. – The species is named after its thick walls (from Latin “*crassus*” = thick, and “*parietis*” = wall).

Holotype. – PMO 221.770.

Paratypes. – PMO 221.750–221.755, PMO 221.757, PMO 221.760, PMO 221.761–221.762, PMO 221.765–221.769, PMO 221.765, PMO 221.807–221.808, PMO 221.810, PMO 221.811, PMO 221.814.

Type locality. – Nes bus stop, Ringerike, Norway.

Type stratum. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian.

Diagnosis. – Branched colonies with distinct wide exozones; autozoecial apertures polygonal; mesozooecia rare; acanthostyles common, spaced 1 to 5 around autozoecial apertures, moderate in size, locally absent; maculae consisting of macrozoecia.

Description. – Branched colonies, branch diameter 2.75–5.20 mm. Exozone distinct, 0.40–1.40 mm wide, endozone 1.55–2.90 mm wide. Secondary overgrowths absent. Autozoecia long, growing parallel to branch axis for a long distance in endozone, in exozone bending, having polygonal shape in transverse section in endozone. Autozoecial apertures polygonal. Diaphragms in endozone rare, planar, thin; in exozone abundant, straight or slightly curved. Mesozooecia rare, small, polygonal in cross section, restricted to exozone. Acanthostyles common, spaced 1 to 5 around autozoecial apertures, moderate in size, having distinct hyaline cores and wide,

Figure 8. *Trematopora maculata* sp. nov. • A–C – longitudinal section, holotype PMO 221.816. • D – tangential section, holotype PMO 221.816. • E – longitudinal section of secondary encrusting, paratype PMO 221.827. • F – longitudinal section, paratype PMO 221.776. • G – transverse section, paratype PMO 221.817. • H – tangential section, paratype PMO 221.824.




Table 6. Measurements of *Amplexopora crassiparietum* sp. nov. Abbreviations as in Table 1.

	N	X	SD	CV	MIN	MAX
Branch width, mm	10	3.91	0.916	23.43	2.75	5.20
Exozone width, mm	10	0.90	0.312	34.64	0.40	1.40
Endozone width, mm	10	2.11	0.439	20.80	1.55	2.90
Aperture width, mm	40	0.11	0.017	15.93	0.08	0.14
Aperture spacing, mm	40	0.20	0.023	11.33	0.15	0.26
Aperture width, mm (macular)	16	0.19	0.030	15.72	0.15	0.25
Aperture spacing, mm (macular)	16	0.26	0.023	9.14	0.22	0.30
Mesozoecia width, mm	30	0.048	0.013	26.91	0.025	0.078
Acanthostyle diameter, mm	30	0.034	0.006	17.36	0.025	0.050
Acanthostyles per aperture	40	3.1	0.888	28.89	1.0	5.0
Autozoecial diaphragms spacing, mm	40	0.15	0.044	29.09	0.07	0.24
Exozonal wall thickness, mm	40	0.079	0.023	29.50	0.030	0.135

Table 7. Measurements of *Amplexopora evae* sp. nov. Abbreviations as in Table 1.

	N	X	SD	CV	MIN	MAX
Branch width, mm	6	3.10	0.334	10.78	2.55	3.40
Exozone width, mm	6	0.74	0.309	41.65	0.35	1.00
Endozone width, mm	6	1.62	0.386	23.82	1.10	2.18
Aperture width, mm	85	0.15	0.026	17.02	0.10	0.21
Aperture spacing, mm	85	0.19	0.031	15.82	0.12	0.30
Aperture width, mm (macular)	16	0.20	0.015	7.35	0.18	0.23
Aperture spacing, mm (macular)	16	0.27	0.039	14.57	0.20	0.33
Acanthostyle diameter, mm	61	0.049	0.013	26.40	0.025	0.080
Acanthostyles per aperture	75	2.7	1.075	39.72	1.0	5.0
Mesozoecia width, mm	60	0.05	0.018	36.31	0.02	0.09
Mesozoecial diaphragms spacing, mm	20	0.12	0.036	30.16	0.07	0.20
Autozoecial diaphragms spacing, mm	49	0.19	0.061	31.70	0.09	0.35
Exozonal wall thickness, mm	40	0.046	0.013	29.05	0.025	0.075

dark sheaths, situated in the dark median lining of autozoecial walls, originating from base of exozone; locally acanthostyles absent. Autozoecial walls in endozone 0.005–0.010 mm thick, granular; in exozone 0.030 to 0.135 mm thick, displaying reverse V-shaped lamination with dark, serrated median lining. Maculae indistinct, 0.8–1.3 mm in diameter, consisting of macrozoecia.

Remarks. – *Amplexopora crassiparietum* sp. nov. differs from *A. silurica* Astrova, 1970 from the lower Silurian (Llandovery) of Ukraine (Podolia) in smaller autozoecial apertures (aperture width 0.08–0.14 mm vs 0.17–0.20 mm in *A. silurica*), and in less abundant acanthostyles (1–5 vs 4–7 per autozoecial aperture in *A. silurica*).

Occurrence. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian; Nes bus stop and Åsa, Ringerike, Norway.

***Amplexopora evae* sp. nov.**

Figures 10A–F, 11A, B, Table 7

Etymology. – The species is named for the junior author’s wife to honour her participation in bryozoan fieldwork.

Holotype. – PMO 170.890.

Paratypes. – PMO 170.886, PMO 170.882, PMO 170.887–170.888, PMO 221.771–221.772, PMO 221.773–221.777, PMO 221.780, PMO 221.804–221.805.

Type locality. – Ødegårdsviken, Ringerike, Norway.

Type stratum. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian.

Diagnosis. – Branched and encrusting colonies; branches with distinct wide exozones; autozoecial apertures polygonal; mesozoecia rare; acanthostyles common, spaced 1 to 5 around autozoecial apertures, moderate in size, locally absent; maculae consisting of macrozoecia.

Description. – Branched colonies, branch diameter 2.55–3.40 mm. Exozone distinct, 0.35–1.00 mm wide, endozone 1.10–2.18 mm wide. Secondary overgrowths common, 0.65–2.50 mm thick. Encrusting colonies 1.5–2.5 mm thick. In branched colonies, autozoecia long, growing parallel to branch axis for a long distance in endozone, in exozone bending, having polygonal shape in transverse section in endozone. In encrusting colonies autozoecia arise directly from the epitheca with narrow endozonal part. Autozoecial apertures polygonal. Diaphragms in endozone rare, planar, thin; in exozone abundant, straight or slightly curved. Mesozoecia rare, small, polygonal in cross section, restricted to exozone. Acanthostyles common, spaced 1 to 5 around autozoecial apertures, moderate in size, having distinct hyaline cores and wide, dark

Figure 9. *Amplexopora crassiparietum* sp. nov. • A – longitudinal section, holotype PMO 221.770. • B, C – transverse section, holotype PMO 221.770. • D – longitudinal section, holotype PMO 221.770. • E – tangential section, holotype PMO 221.770. • F – tangential section, paratype PMO 221.760.




Table 8. Measurements of *Eridotrypella* sp. Abbreviations as in Table 1.

	N	X	SD	CV	MIN	MAX
Aperture width, mm	20	0.11	0.013	11.15	0.08	0.13
Aperture spacing, mm	20	0.19	0.022	11.51	0.16	0.22
Exilazooecia width, mm	7	0.05	0.014	29.91	0.03	0.07

sheaths, situated in the dark median lining of autozooeical walls, originating from base of exozone; locally acanthostyles absent. Autozooeical walls in endozone 0.005–0.010 mm thick, granular; in exozone 0.025–0.075 mm thick, displaying reverse V-shaped lamination with dark, serrated median lining. Maculae indistinct, consisting of macrozoecia, 0.70–1.35 mm in diameter.

Remarks. – *Amplexopora evae* sp. nov. differs from *A. crassiparietum* sp. nov. in thinner autozooeical walls and larger autozooeical apertures.

Occurrence. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian; Ødegårdsviken, Ringerike, Norway.

Family Eridotrypellidae Morozova, 1961

Genus *Eridotrypella* Duncan, 1939

Type species. – *Batostomella obliqua* Ulrich, 1890. Middle Devonian; Michigan (USA).

Diagnosis. – Branched colonies. Autozooeical apertures irregularly polygonal. Autozooeical walls laminated, without distinct zooeical boundaries, irregularly thickened, containing spherulites. Diaphragms complete, varying in number. Exilazooecia rare. Acanthostyles varying in size and number.

Comparison. – *Eridotrypella* Duncan, 1939 differs from *Eostenopora* Duncan, 1939 in colony form (ramose branched vs encrusting or massive colonies).

Occurrence. – Silurian–?Carboniferous; worldwide.

***Eridotrypella* sp. A**

Figure 11C–H, Table 8

Material. – PMO 221.789, PMO 221.782, PMO 221.796.

Description. – Ramose branched colonies, 1.2 mm in diameter. Exozone 0.34 mm wide, endozone 0.52 mm wide. Exozones distinctly separated from endozones. Secondary overgrowth occurring, 0.28–0.66 mm thick. Autozooeica long in endozones, bending sharply in exozones. Autozooeical apertures polygonal with rounded corners. Autozooeical diaphragms absent to rare in endozones; common to abundant in transition between endozone and exozone, straight or inclined. Exilazooecia rare, short, polygonal in cross section. Acanthostyles rare, having distinct narrow cores and laminated sheaths, 0.030–0.035 mm in diameter. Autozooeical walls granular, locally weakly crenulated, 0.005–0.010 mm thick in endozones; serrated in the longitudinal view and merged in the tangential section, 0.04–0.06 mm thick in exozones. Maculae not observed.

Remarks. – *Eridotrypella* sp. differs from *E. sepizensis* Astrova, 1970 from the lower Silurian (Wenlock) of Estonia in less abundant acanthostyles and in larger apertures (aperture width 0.08–0.13 mm vs 0.13–0.20 mm in *E. sepizensis*). *Eridotrypella* sp. differs from *E. duncaniae* Pushkin, 1976 from the upper Silurian (Ludlow) of Belarus in narrower branches (branch width 1.2 mm vs 2.5–5.0 mm in *E. duncaniae*), and in smaller apertures (aperture width 0.08–0.13 mm vs 0.14–0.25 mm in *E. duncaniae*).

Occurrence. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian; Ødegårdsviken, Ringerike, Norway.

Order Cryptostomata Vine, 1884

Suborder Rhabdomesina Astrova & Morozova, 1956

Family Rhabdomesidae Vine, 1884

Genus *Orthopora* Hall, 1886

Type species. – *Trematopora regularis* Hall, 1874. Lower Devonian; USA.

Diagnosis. – Branched colonies. Autozooeica short, budding from more or less distinct medial axis in spiral order. Autozooeical diaphragms rare to absent. Both superior and inferior hemisepta commonly present; sometimes double hemisepta occurring; rarely hemisepta absent. Autozooeical apertures oval, arranged regularly in alternating rows on the colony surface. Walls granular in the endozone; laminated in exozone. Paurostyles abundant, prominent. Acanthostyles present, less abundant than paurostyles. Heterozooecia absent.

Figure 10. *Amplexopora evae* sp. nov., holotype PMO 170.890. • A – general view of a branched colony. • B – transverse section. • C – longitudinal section. • D – longitudinal section of encrusting colony. • E – transverse section showing wall structure. • F – tangential section.




Table 9. Measurements of *Orthopora worsleyi* sp. nov. Abbreviations as in Table 1.

	N	X	SD	CV	MIN	MAX
Branch width, mm	21	0.59	0.098	16.57	0.42	0.77
Exozone width, mm	21	0.17	0.041	24.03	0.10	0.25
Endozone width, mm	21	0.25	0.064	25.33	0.13	0.35
Aperture width, mm	45	0.06	0.012	19.06	0.04	0.08
Aperture spacing along branch, mm	25	0.31	0.045	14.63	0.21	0.40
Aperture spacing diagonally, mm	25	0.15	0.019	12.57	0.12	0.20
Acanthostyle diameter, mm	30	0.031	0.006	18.38	0.025	0.045

Table 10. Measurements of *Mediaporina kiaeri* sp. nov. Abbreviations as in Table 1.

	N	X	SD	CV	MIN	MAX
Aperture width, mm	19	0.08	0.014	17.79	0.06	0.10
Aperture spacing, mm	20	0.17	0.017	9.59	0.14	0.2
Heterozooecia width, mm	20	0.03	0.008	28.09	0.015	0.045
Acanthostyle diameter, mm	20	0.034	0.006	19.03	0.025	0.050

Remarks. – *Orthopora* Hall, 1886 differs from *Trematella* Hall, 1886 in absence of metazooecia and in presence of well-developed hemisepta.

Occurrence. – Silurian to Carboniferous of North America, Europe and China, Middle Permian of Oman.

***Orthopora worsleyi* sp. nov.**

Figure 13A–H, Table 9

Etymology. – The species is named for David Worsley, who has been an inspiration to us in the study of Silurian rocks of the Oslo Region.

Holotype. – PMO 221.790.

Paratypes. – PMO 221.750–221.755, PMO 221.757, PMO 221.781–221.782, PMO 221.784–221.785, PMO 221.789, PMO 221.794, PMO 221.796.

Type locality. – Nes bus stop, Ringerike Norway.

Type stratum. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian.

Diagnosis. – Branched colonies; bifurcation common; both superior and inferior hemisepta well developed; acanthostyles abundant, arranged in single row between autozooecia, varying in size; paurostyles abundant.

Description. – Branched colonies arising from encrusting bases. Branches 0.42–0.77 mm in diameter, with 0.13–0.35 mm wide endozones and 0.10–0.25 mm wide exozones. Branch bifurcation common. Transverse sections of branches circular. Autozooecia medium in length, budding from indistinct medial axis in spiral order. Autozooecial diaphragms rare to absent. Superior hemisepta present, blunt, curved proximally, positioned near the bend of autozooecial chamber; inferior hemisepta long, positioned beneath superior hemisepta, curved distally. Autozooecial apertures oval, arranged regularly in alternating rows on the colony surface. Walls in the endozone granular, 0.003–0.005 mm thick; laminated in exozone. Acanthostyles abundant, arranged in longitudinal rows between apertures, varying in size, having narrow hyaline cores and wide laminated sheaths. Paurostyles abundant, arranged irregularly between acanthostyles, 0.008–0.016 mm in diameter. Heterozooecia absent.

Remarks. – *Orthopora worsleyi* sp. nov. differs from *O. casualis* Goryunova, 1985 from the lower Silurian Rochester Shale (Wenlock) of New York, USA in narrower branches (0.42–0.77 mm vs 0.80–1.20 mm in *O. casualis*), and in slightly smaller apertures (aperture width 0.04–0.08 mm vs 0.05–0.10 mm in *O. casualis*). *Orthopora worsleyi* sp. nov. differs from species described by Kopaeovich (1975) as *Orthopora rhombifera* (Hall, 1874) from the upper Silurian (Přídolí) of Estonia in narrower branches (0.42–0.77 mm vs 0.76–1.17 mm in that species).

Occurrence. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian; Ødegårdsviken, Ringerike, Norway.


Family Arthrostylidae Ulrich, 1882

***Mediaporina Pushkin* in Pushkin et al., 1990**

Type species. – *Mediapora aspersa* Kopaeovich, 1975. Upper Silurian, Ludlow; Estonia.

Diagnosis. – Branched colonies. Autozooecia relatively

Figure 11. A, B – *Amplexopora evae* sp. nov., tangential section, holotype PMO 170.890. • C–H – *Eridotrypella* sp.; C, D – transverse section, paratype PMO 221.782. • E – longitudinal section, holotype PMO 221.789. • F, G – tangential section, paratype PMO 221.789. • H – longitudinal section, holotype PMO 221.789.






Figure 12. *Mediaporina kiaeri* sp. nov. • A, B – longitudinal section, holotype PMO 221.754. • C – oblique section, paratype PMO 221.762. • D, E – tangential section showing autozoecial apertures, styles and heterozoecia, holotype PMO 221.754. • F – deep tangential section showing autozoecial apertures and heterozoecia, paratype PMO 221.762. • G – longitudinal section showing autozoecia and heterozoecia, paratype PMO 221.762.

short, budding from more or less distinct medial axis in spiral order. Autozoecial diaphragms rare to common. Hemisepta absent. Autozoecial apertures oval, arranged regularly in alternating rows on the colony surface. Walls granular in the endozone; laminated in exozone. Paurostyles large, abundant, prominent. Acanthostyles absent. Heterozoecia tubular, common to abundant, restricted to exozone, bearing few diaphragms.

Remarks. – *Mediaporina* Pushkin in Pushkin *et al.*, 1990 differs from *Sceptropora* Ulrich, 1888 in having branched

colony instead of branched segments with expanded distal parts.


Occurrence. – Lower to upper Silurian of Europe (British Isles, Norway, Estonia, Belarus, Latvia).

***Mediaporina kiaeri* sp. nov.**

Figure 12A–G, Table 10

Etymology. – The species name refers to Johan A. Kiær

Figure 13. *Orthopora worsleyi* sp. nov. • A, B – oblique section, holotype PMO 221.790. • C, D – oblique section of the basal part of the colony, paratype PMO 221.794. • E – oblique section of dichotomous colony, PMO 221.783. • F, G – longitudinal section, PMO 221.784. • H – branch transverse section, paratype PMO 221.783.



who described the Silurian rocks of Ringerike in great detail in 1908.

Holotype. – PMO 221.754.

Paratype. – PMO 221.762.

Type locality. – Nes bus stop, Ringerike Norway.

Type stratum. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian.

Diagnosis. – Branched colonies; autozooeical diaphragms rare to common; acanthostyles abundant, 7–9 surrounding each autozooeica aperture; heterozooeica abundant, 4–6 surrounding each autozooeica aperture.

Description. – Branches 1.05–1.20 mm in diameter, with 0.30–0.37 mm wide endozones and 0.34–0.45 mm wide exozones. Branch bifurcation not observed. Autozooeica moderately long, growing along the branch axis in endozone and bending sharply in exozone, polygonal in transverse section of endozone. Autozooeical diaphragms rare to common, 2–3 usually present in each autozooeicum. Hemisepta absent. Autozooeical apertures oval to slightly angular, arranged regularly in alternating rows on the colony surface. Walls in the endozone granular, 0.003–0.005 mm thick; indistinctly laminated in exozone. Acanthostyles abundant, 7–9 surrounding each autozooeica aperture, restricted to exozone, varying in size, having indistinct wide cores and narrow sheaths. Heterozooeica abundant, 4–6 surrounding each autozooeica aperture, originating at the base of exozone, rounded to oval at colony surface, containing thin diaphragms, often sealed by calcitic material at colony surface.

Remarks. – *Mediaporina kiaeri* sp. nov. differs from *M. aspersa* (Kopaeovich, 1975) from the upper Silurian of Estonia by more abundant and smaller heterozooeica (heterozooeica width 0.015–0.045 mm vs 0.05–0.13 mm in *M. aspersa*) and by larger styles (style diameter 0.025–0.050 mm vs 0.02–0.04 mm in *M. aspersa*). *Mediaporina kiaeri* sp. nov. differs from *M. orbiculata* (Pushkin, 1975) from the upper Silurian of Belarus in smaller autozooeical apertures (aperture width 0.06–0.10 mm vs 0.09–0.12 mm in *M. orbiculata*) and by more abundant heterozooeica.

Occurrence. – Steinsfjorden Formation, Brattstad Member, Silurian, Wenlock, Sheinwoodian–Homerian; Nes bus stop, Ringerike, Norway.

Acknowledgements

The Deutsche Akademische Austauschdienst (DAAD) supported Andrej Ernst's study visit to the Natural History Museum, Uni-

versity of Oslo, with a one-year fellowship 2002–2003 (grant D/02/00949). Patrick N. Wyse Jackson, Dublin, and Caroline Buttler, Cardiff, are thanked for their helpful reviews.

References

- ANSTEY, R.L. & PERRY, T.G. 1970. Biometric procedures in taxonomic studies of Paleozoic bryozoans. *Journal of Paleontology* 44, 383–398.
- ASTROVA, G.G. 1957. Some new species of bryozoans from the Silurian of Tuva. *Materiali k osnovam paleontologii Akademii nauk SSSR* 1, 5–14. [in Russian]
- ASTROVA, G.G. 1959. Silurian bryozoans of the central and western Tuva. *Trudy Paleontologicheskogo instituta Akademii nauk SSSR* 79, 1–74. [in Russian]
- ASTROVA, G.G. 1964. New Order of the Paleozoic Bryozoa. *Paleontologicheskii zhurnal* 1964(2), 22–31. [in Russian]
- ASTROVA, G.G. 1965. Morphology, history of development and systematics of the Ordovician and Silurian Bryozoa. *Trudy Paleontologicheskogo instituta Akademii nauk SSSR* 106, 1–432. [in Russian]
- ASTROVA, G.G. 1970. New Silurian and Early Devonian bryozoans of Orders Cystoporata and Trepostomata from Estonia and Podolia, 7–22. In ASTROVA, G.G. & CHUDINOVA, I.I. (eds) *New species of Palaeozoic bryozoans and corals*. Nauka, Moscow.
- ASTROVA, G.G. 1978. The history of development, system, and phylogeny of the Bryozoa: Order Trepostomata. *Trudy Paleontologicheskogo instituta Akademii nauk SSSR* 169, 1–240. [in Russian]
- ASTROVA, G.G. & MOROZOVA, I.P. 1956. On systematics of the order Cryptostomata. *Doklady Akademii nauk SSSR* 110(4), 661–664. [in Russian]
- BASSLER, R.S. 1906. The bryozoan fauna of the Rochester Shale. *U.S. Geological Survey Bulletin* 292, 1–137.
- BOARDMAN, R.S. 1960. A revision of the Ordovician bryozoan genera *Batostoma*, *Anaphragma*, and *Amplexopora*. *Smithsonian Miscellaneous Collections* 140(5), 1–28.
- BORG, F. 1926. Studies on Recent cyclostomatous Bryozoa. *Zoologiska Bidrag från Uppsala* 10, 181–507.
- DUNCAN, H. 1939. Trepostomatous Bryozoa from the Traverse Group of Michigan. *University of Michigan Paleontology Contributions* 5(10), 171–270.
- EHRENBERG, C.G. 1831. *Symbolae Physicae, seu Icones et descriptiones Corporum Naturalium novorum aut minus cognitorum, quae ex itineribus per Libyam, Aegyptum, Nubiam, Dongalaam, Syriam, Arabiam et Habessiniam, studia annis 1820–25, redirent. Pars Zoologica, 4, Animalia Evertibrata exclusis Insectis*. 10 pls. Berolini.
- GORYUNOVA, R.V. 1985. Morphology, system and phylogeny of Bryozoa (Order Rhabdomesida). *Trudy Paleontologicheskogo instituta Akademii nauk SSSR* 208, 1–152. [in Russian]
- HAGEMAN, S.J. 1993. Effects of nonnormality on studies of the morphological variation of a rhabdomesine bryozoan, *Streblotrypa* (*Streblascopora*) *prisca* (Gabb & Horn). *The University of Kansas Paleontological Contributions* 4, 1–13.

- HALL, J. 1852. Organic remains of the lower middle division of the New York system. *Natural History of New York. Part 6. Palaeontology of New York* 2, 40–52, 144–173.
- HALL, J. 1874. Descriptions of Bryozoa and corals of the lower Helderberg Group. *Twenty-sixth annual report of the New York State Museum of Natural History*, 93–115.
- HALL, J. 1886. Bryozoa of the Upper Helderberg Group; plates and explanations. *State Geologist of New York, Annual Report for 1885*, 25–53.
- HENNIG, A. 1908. Gotlands Silurbryozoen 3. *Arkiv för zoologi* 4, 1–64.
- HETHERINGTON, C.J., NAKREM, H.A. & POTEI, S. 2011. Note on the composition and mineralogy of Wenlock Silurian bentonites from the Ringerike District: Implications for local and regional stratigraphic correlation and sedimentary environments. *Norwegian Journal of Geology* 91, 181–192.
- KLIER, J. 1908. *Das Obersilur im Kristianitgebiete. Eine stratigraphisch-faunistische Untersuchung. Skrifter udgivne af Videnskabs-selskabet i Christiania*. 595 pp. J. Dybwad, Christiania.
- KOPAEVICH, G.V. 1975. Silurian Bryozoa of Estonia and Podolia (Cryptostomata and Rhabdomesonata). *Trudy Paleontologicheskogo instituta Akademii nauk SSSR* 151, 5–153. [in Russian]
- KOPAEVICH, G.V. 1984. Atlas of Bryozoa from the Ordovician, Silurian and Devonian of Mongolia. *Sovmestnaya sovetko-mongolskaya paleontologicheskaya ekspeditsiya, Trudy* 22, 1–164. [in Russian]
- M'COY, F. 1849. On some new genera and species of Palaeozoic corals and Foraminifera. *Annals and Magazine of Natural History* 3(2), 119–136.
DOI 10.1080/03745485909494606
- MELCHIN, M.J., SADLER, P.M. & CRAMER, B.D. 2012. Chapter 21: The Silurian Period, 525–558. In GRADSTEIN, F.M., OGG, J.G., SCHMITZ, M. & OGG, G. (eds) *The Geologic Time Scale 2012*. Elsevier, New York.
- MILLER, S.A. 1889. *North American Geology and Paleontology for the use of amateurs, students, and scientists*. 664 pp. Western Methodist Book concern, Cincinnati.
- MILNE-EDWARDS, H. & HAIME, J. 1851. Monographie des poly-piers fossiles des terrains paléozoïques, précédé d'un tableau général de la classification des Polypes. *Archives du Muséum d'histoire naturelle* 5, 1–504.
- MOROZOVA, I.P. 1961. Devonian Bryozoa of the Minusinsk and Kuznetsk basins. *Trudy Paleontologicheskogo instituta Akademii nauk SSSR* 86, 1–207. [in Russian]
- NICHOLSON, H.A. 1879. *On the structure and affinities of the "Tabulate Corals" of the Paleozoic period, with critical descriptions of illustrative species*. 342 pp. William Blackwood and Sons, Edinburgh.
- NICHOLSON, H.A. & FOORD, A.H. 1885. On the genus *Fistulipora* M' Coy, with descriptions of several species. *Annals and Magazine of Natural History, Series 5* 16(96), 496–517.
DOI 10.1080/00222938509459918
- OLAUSSEN, S. 1981. Formation of celestite in the Wenlock, Oslo Region, Norway – evidence for evaporitic depositional environments. *Journal of Sedimentary Petrology* 51, 37–46.
- ORBIGNY, A.D. D' 1850. *Prodrome de paléontologie stratigraphique universelle des animaux mollusques rayonnés, faisant suite ou cours élémentaire de paléontologie et géologie stratigraphique. Volume 1*. 394 pp. Victor Masson, Paris.
DOI 10.5962/bhl.title.62810
- OWEN, D. E. 1969. Wenlockian bryozoa from Dudley, Niagara, and Gotland and their palaeogeographic implications. *Palaeontology* 12(4), 621–636.
- PUSHKIN, V.I. 1975. About genus *Mediapora* and its new representatives from the Upper Silurian in Belarus and Latvia. *Transactions of the Museum of earth-study of Moscow University* 11, 113–120. [in Russian]
- PUSHKIN, V.I. 1976. New species of Ordovician and Silurian Bryozoa of the Brest Depression, 3–40. In AKIMETS, V.S. & GOVETSKII, G.I. (eds) *New species of the fossil plants and animals of Belarus*. Nauka i tekhnika, Minsk. [in Russian]
- PUSHKIN, V.I., NEKHOROSHEVA, L.V., KOPAEVICH, G.V. & YAROSHINSKAYA, A.M. 1990. *Przhydolian Bryozoa of the USSR*. 125 pp. Nauka, Moscow. [in Russian]
- SPIJELDNÆS, N. 1982. Silurian Bryozoa of the Oslo Region. IUGS Subcommittee on Silurian Stratigraphy. Field Meeting, Oslo Region 1982. *Paleontological Contributions from the University of Oslo* 278, 55–62.
- ULRICH, E.O. 1879. Description of a new genus and some new species of bryozoans from the Cincinnati Group. *Journal of the Cincinnati Society of Natural History* 2, 119–131.
- ULRICH, E.O. 1882. American Palaeozoic Bryozoa. *The Journal of the Cincinnati Society of Natural History* 5, 121–175, 233–257.
- ULRICH, E.O. 1888. On *Sceptropora*, a new genus of Bryozoa, with some remarks on *Helopora* Hall, and other genera of that type. *The American Geologist* 1(4), 228–234.
- ULRICH, E.O. 1890. Palaeozoic Bryozoa, 283–688. In WORTHEN, A.H. & LINDAHL, J. (eds) *Geological Survey of Illinois, Vol. 8, Geology and Palaeontology, Part II*. Geological Survey of Illinois, Springfield.
- ULRICH, E.O. 1893. On Lower Silurian Bryozoa of Minnesota. *The Geological and Natural History Survey of Minnesota, final report* 3, 96–332.
- VINE, G.R. 1884. Fourth report of the Committee consisting of Dr. H.R. Sorby and Mr. G.R. Vine, appointed for the purpose of reporting on fossil Polyzoa, 161–209. *Report of the Fifty-third meeting of the British Association for the Advancement of Science (Southport, 1883)*. John Murray, London.
- WHITAKER, J.H.M. 1977. *A Guide to the Geology around Steinsfjord, Ringerike*. 56 pp. Universitetsforlaget, Oslo.
- WORSLEY, D., AARHUS, N., BASSETT, M.G., HOWE, M.P.A., MØRK, A. & OLAUSSEN, S. 1983. The Silurian succession of the Oslo Region. *Norges geologiske undersøkelse* 384, 1–57.
- WORSLEY, D., BAARLI, B.G., HOWE, M.P.A., HJALTASON, F. & ALM, D. 2011. New data on the Bruflat Formation and the Llandoverly/Wenlock transition in the Oslo Region, Norway. *Norwegian Journal of Geology* 91, 101–120.